ESRD Special Project

Developing Clinical Performance Measures for the Care of Patients with End Stage Renal Disease

Phase II:

2000 Study of ESRD Clinical Performance Measures: Reliability Report

Deliverable #23 Reliability Testing Report on the 2000 CPMs Data Collection

January 19, 2001

Submitted to:

Mary Beth Ribar Acute & Chronic Disease Management Division QMHAG, OCSQ Health Care Financing Administration 7500 Security Boulevard, S3-02-01 Baltimore, MD 21244 Nancy Tollison, Project Officer Health Care Financing Administration Region X 2201 6th Avenue, MS/RX-42 Seattle, WA 98121

Submitted by:

Jonathan Sugarman, MD, MPH, Project Director Naomi Maxey, Project Coordinator Lisa Zhao, MS, Data Analyst PRO-West, A Professional Review Organization 10700 Meridian Avenue North, Suite 100 Seattle, WA 98133



Table of Contents

Contents

Intr	oduction	1
Bac	ckground	1
Pro	eject Methods	1
	Statistical Methods	
	Concurrence	2
	Kappa Statistics	2
	T-Test statistics Based on Means	2
	Data Collection	2
	Sample and Measures: Hemodialysis CPMs	3
	Sample and Measures: Peritoneal dialysis CPMs	3
	Sample and Measures: Hemodialysis Facility Forms	4
Res	sults	4
	Hemodialysis CPMs.	4
	Peritoneal Dialysis CPMs	5
	Hemodialysis Facility Forms	5
Cor	nclusions	6
Table	es	
	Table 1: Calculation of data concurrence	7
	Table 2: Comparison of categorical data abstracted by dialysis facility staff to categorical data re-abstracted by ESRD Network staff for selected hemodialysis measures (excluding measures related to vascular access)	8

fable 3: Comparison of means for continuous data abstracted by dialysis facility staff to continuous data re-abstracted by ESRD Network staff for selected hemodialysis measures (excluding measures related to vascular access)
Table 4: Reported Kt/V [Question16I]
Table 5: Method used to calculate the reported Kt/V [Question 16J]
Table 6: Hemoglobin ≥ 9gm/dL [Question 15A]
Table 7: Hemoglobin ≥ 11gm/dL [Question 15A]
Table 8: Transferrin saturation $\geq 20\%$ [Question 15F]
Table 9: Serum ferritin concentration ≥ 100 ng/dL [Question 15E]
Table 10: Epoetin dosage prescription (Y/N) [Question 15B]
Table 11: Prescribed route of epoetin administration [Question 15D]
Table 12: Serum albumin values (≥ 3.5/3.2 gm/dL by BCG/BCP methods) [Question 17A]
Table 13: Laboratory method used to measure serum albumin in Table 12 [Question 17B]
Table 14: The type of access used on the last hemodialysis session on or between October 1, 1999, and December 31, 1999 [Question 18A]
Table 15: Reason for catheter, if used for access between October 1, 1999, and December 31, 1999 [Question 18B1]
Table 16: Catheter duration (\$90 days), if used for access between October 1, 1999, and December 31, 1999 [Question 18B2]
Table 17: Presence of routine monitoring for stenosis, when synthetic grafts, bovine grafts, or AV fistulas were used for access between October 1, 1999, and December 31, 1999 [Question 18C1]
Table 17a-e: The routine stenosis monitoring method used between October 1, 1999, and December 31, 1999, when synthetic grafts, bovine grafts, or AV fistulas were used for access [Question 18C2]
Table 18: The type of access used at the initiation (re-initiation) of the first time hemodialysis, if between January 1, 1999 – August 31, 1999 [Question 19A]

Table 19: The type of access used 90 days after the date in Table 18 during the initiation (re-initiation) of hemodialysis, if between	
January 1, 1999 – August 31, 1999 [Question 19C]	21
Table 20: Limb amputation (Y/N) [Question 13]	21
Table 21: Number of scheduled hemodialysis times per week [Question 16A]	22
Table 22: Agreement rate of data abstracted by dialysis facility staff to data re-abstracted by ESRD Network staff for selected hemodialysis measures	22
Table 23: Comparison of categorical data abstracted by dialysis facility staff to categorical data re-abstracted by ESRD Network staff for selected peritoneal dialysis measures	23
Table 24: Comparison of means for continuous data abstracted by dialysis facility staff to continuous data re-abstracted by ESRD Network staff for selected peritoneal dialysis measures.	24
Table 25: Reported Kt/V [Question 18D]	25
Table 26: Method used to calculate the V in Kt/V _{urea} [Question 18E]	25
Table 27: Reported weekly creatinine clearance [Question 18F]	26
Table 28: Creatinine clearance corrected for body surface area, using standard methods (Y/N) [Question 18G]	26
Table 29: Hemoglobin ≥ 9 gm/dL [Question 15A]	26
Table 30: Hemoglobin ≥ 11 gm/dL [Question 15A]	26
Table 31: Transferrin saturation ≥ 20% [Question 15F]	27
Table 32: Serum ferritin concentration ≥ 100 g/mL [Question 15E]	27
Table 33: Epoetin dose prescription (Y/N) [Question 15B]	27
Table 34: Prescribed route of epoetin administration [Question 15D]	27
Table 35: Serum albumin values (≥ 3.5/3.2 gm/dL by BCG/BCP methods) [Question 16A]	28
Table 36: Laboratory method used to measure serum albumin in Table 35 [Question 16B]	28

Table 37: Number of scheduled dialysis days per week [Question 19A]
Table 38: Total number of exchanges per 24 hours for CAPD patients [Question 19B2]
Table 39: Total number of dialysis exchanges during the nighttime for cycler patients [Question 19C3b]
Table 40: Total number of dialysis exchanges during the daytime for cycler patients [Question 19C4b]
Table 41: Prescription changed (Y/N) [Question 19E2]
Table 42: Limb amputation (Y/N) [Question 13]
Table 43: Agreement rate of data abstracted by dialysis facility staff to data re-abstracted by ESRD Network staff for selected peritoneal dialysis measures
Table 44: Do facilities have a written policy for the timing of the post-BUN sample collection? [Question 1]
Table 45: Facilities' written policy regarding post-dialysis BUN sampling [Question 1]
Table 46: Facilities' compliance with written policy (internal audit) regarding post-dialysis BUN sampling [Question 2]
Table 47: Facilities that re-used dialyzers from October 1, 1998, to
December 31, 1998 [Question 3]
Table 48: Facilities that pre-volumed 95-100% of dialyzers intended for reuse [Question 3]
Table 49: Facilities that pre-volumed <95% of dialyzers intended for reuse [Question 3]
Table 50: Facilities that used batch testing and/or of an average TCV for a
group of hemodialyzers to infer TCV [Question 3]
Table 51: Facilities that used the manufacturer's product information to infer TCV [Question 3]